



Fungal endophytes of an aquatic weed *Marsilea minuta* Linn

Udaya Prakash NK¹, Ashwin Karthick N², Poomagal D², Susithra M², Chandran M³ and Bhuvaneshwari S⁴

¹ Department of Biotechnology, School of Life Sciences, Vels University, Pallavaram, Chennai, India.

² R and D, MARINA LABS, Nerkundram, Chennai, India.

³ Department of Biotechnology, VelTech HighTech Dr. RR Dr. SR Engineering College, Avadi, India

⁴ Department of Botany, Bharathi Women's College, Chennai, India.

Udaya Prakash NK, Ashwin Karthick N, Poomagal D, Susithra M, Chandran M, Bhuvaneshwari S 2018 – Fungal endophytes of an aquatic weed *Marsilea minuta* Linn. Current Research in Environmental & Applied Mycology 8(1), 86–95, Doi 10.5943/cream/8/1/7

Abstract

Endophytic fungi are the organisms that colonize a plant without causing apparent harm to the host at any time in their life cycle. Endophytes from different angiosperms are widely studied however such studies from Pteridophytes are rare. In this study, the plant *Marsilea minuta*, an aquatic Pteridophyte belonging to the family Marsileaceae was explored for the presence of endophytic fungi. The plant was collected from four different places in and around Chennai (Avadi, Ambattur, Chengalpet and Guduvanchery). A total of 800 segment consisting leaflets, stolens, runners and roots (200 segments each) were investigated for the presence of endophytes. The study resulted in isolation of 574 colonies belonging to 17 fungal species among which 14 forms the Hyphomycetes, three forms the Coelomycetes and one form the non sporulating morphospecies. The data obtained were analysed statistically for relative density, colonization frequency, Species diversity indices like Gleason index, Shannon index, and Simpson's dominance index.

Keywords – Aquatic – Endophytic Fungi – Pteridophyte – Statistical analysis

Introduction

The term endophyte is defined by many authors, commonly as non-pathogenic organisms living within plant cell without causing disease symptoms or external structural alterations (Azevedo & Araujo 2007, Hyde & Soyong 2008, Sakalidis et al. 2011). Hirsch & Braun (1992) stated that the endophytes colonizing living plant tissues don't cause any immediate or overt negative impact. The role of endophytic fungi in ecological systems is varied and diverse. They support the host plant in drought tolerance and vigor (Stone et al. 2004). As mutualists they protect the plant against preying insects (Carroll 1991) and pests (Rocha et al. 2011). Further they help protect the host plants against oxidative stress by stimulating or enhancing the production of antioxidants (White & Torres 2010). Novel, biologically active metabolites of the host plant could be produced in large quantities from the associated endophytic fungi (Selim et al. 2011, Qadri et al. 2014, Tiwari et al. 2014). Thus, endophyte fungi could serve as a potential source for production of these metabolites and for isolation of new drugs (Kumar et al. 2005). Endophytes are ubiquitous in various plants (Arnold & Lutzoni 2007, U'Ren et al. 2012) which includes aquatic plants (Li et al. 2010). Numerous studies have been carried out on the diversity of endophytes in vascular tissues of

higher plants of terrestrial nature due to their diverse ecological roles. However, the endophytic diversity of aquatic Pteridophytes has been rarely studied.

Marsilea minuta L., an aquatic Pteridophyte belonging to the family Marsileaceae, is a small creeping fern with erect tetrafoliate leaves (Parrotta 2001). It is found throughout Africa, Madagascar and Comoros with wide occurrence in tropical Asia and as a weed in the southern United States. The bright green leaves are tender and are eaten as potherb in Senegal, Gambia and India. The weed is known to have traditional medicinal value and is used to stop nose bleeding, treat indigestion. The whole plant is used as sweet, astringent, coolant and expectorant. It is used in the treatment of psychopathy, diarrhoea, cough, skin diseases and fever as reported in Ayurveda. The reported medicinal properties of the plant includes anti- hepatitis and as anti-diabetic (Madhu et al. 2012). Many bioactivities of *M. minuta* have been identified (Parihar et al. 2003, Bhattamisra et al. 2007). Thus the present study aims to explore the endophytic population of the different plant parts of *Marsilea minuta* and further to study the fungal species diversity associated with geographical variation.

Materials & Methods

Sample collection

The healthy plants of *Marsilea minuta* were collected from four different places in and around Chennai (Avadi, Ambattur, Chengalpet and Guduvanchery) in sterile polythene bags and were processed within 24 hours.

Isolation of endophytes

The collected samples were washed thoroughly in running tap water. From the plant *Marsilea minuta*, 50 segments of different tissues of leaflet, petiole, rhizome and root each was cut into approximately 1cm² segments were surface sterilised following the method of Dobranic et al. (1995). The segments were washed in 1% mercuric chloride for 10 seconds, sequentially in 70% ethanol for 30 seconds and finally rinsed with sterile water for 5 seconds. Ten segments of each plant tissues of different geographical locations were placed on Potato Dextrose Agar (PDA) medium amended with streptomycin.

The petri dishes were incubated in a light chamber for a period of three weeks at 28±2°C temperature by maintaining dark -light cycle (Suryanarayanan 1992). The light chamber had a bank of three 4-foot Philips day light fluorescent lamps. The segments received 2200 lux of light through the Petri dishes lid as measured by a Lutron (Germany) 1 × -10¹ Lux meter. The incubation temperature was 28 ± 2° C. The petri dishes were observed periodically and the growing fungi were transferred to fresh PDA slants. To prevent the rapidly growing fungi from inhibiting the slow growing species, the former were removed following isolation (Bills 1996). Sporulating isolates were identified to species level based on their microscopic and macroscopic morphology using standard manuals (Ellis 1971, 1976, Sutton 1980, Udayaprakash 2004). The sterile isolates which could not be assigned to any taxonomic group were sorted described as 'sterile form' in analysis of the results (Frohlich et al. 2000, Suryanarayanan et al. 2000)

Statistical Analysis

The density of colonization (rD %) of single endophytes species and the colonization frequency of endophytic fungi in the segments of *Marsilea minuta* were calculated by the method of Fisher & Petrini (1987).

$$\text{Relative density of Colonization} = \frac{\text{Total no of individuals of a fungi recorded}}{\text{Total number of segments screened}} * 100$$

$$\text{Colonization frequency} = \frac{\text{Number of segments colonized with } \geq 1 \text{ isolate}}{\text{Total number of segments screened}} * 100$$

Species diversity indices

The species diversity indices were calculated in accordance to Kumar & Hyde (2004)

$$\text{Gleason index } [H_G] = N_{p-1}/\ln N_i$$

Where N_i is the total number of fungal isolates, N_p is the total number of species to which these fungi belong

$$\text{Shannon index } [H_S] = -\sum_j (p_j \ln p_j), j = 1, \dots, N_p,$$

Where j is the order given to the individual fungal species, p_j is the proportion of individuals belonging to a particular species (i.e. j^{th} species) to the total number of individuals isolated.

$$\text{Simpson dominance index } (D) = 1/\sum_j (p_j^2)$$

Jaccard's Similarity index

Jaccard's Similarity Coefficients were calculated for all possible pairs of host tissues to compare the endophytic assemblages according to the formula:

$$\text{Similarity coefficient} = \frac{C}{A+B-C}$$

where, A and B are the total number of fungal species isolated from any two tissue type and C the number of fungal species found in common between them (Arnold et al. 2000).

Results

Endophytic fungi composition

The segments of leaflet, petiole, rhizome and root of *Marsilea minuta* were processed for isolation of endophytic fungi from the plant samples collected from four different places in and around Chennai (Avadi, Ambattur, Chengalpet and Guduvanchery). A total of 574 isolates were isolated from 800 segments of the plant. Altogether 17 fungal species were identified based on the morphology of the fungal culture and characteristics of the spore. Among the fungal species fourteen forms belongs to hyphomycetes and three forms to coelomycetes. The non-sporulating cultures were grouped together as sterile forms. The sporulating species accounted for 91.99% of the total isolates, while non-sporulating contributed 8.01%.

The leaflet segments were harboured with 158 isolates belonging to 13 species. The leaflet segments were dominated by the species *Colletotrichum* sp. (43.03%) followed by *Curvularia lunata* (25.31%) and *Drechslera halodes* (16.45%). *Curvularia lunata* was isolated from leaflet segments of Ambattur, Chengalpet and Kovipathagai, while the occurrence of *Colletotrichum* sp. was constrained to the leaflet segments from Guduvanchery and *Drechslera halodes* to leaflet segments of Kovipathagai site.

Segments of the petiole resulted in the isolation of 262 colonies belonging to 8 species. The petiole segments colonized in high proportion by *Curvularia lunata* (38.18%) followed by *Colletotrichum* sp. (18.32%) and *Drechslera halodes* (13.73%). Similar to leaflet segments, the occurrence of *Curvularia lunata* was high in petiole segments of Ambattur, Chengalpet and Kovipathagai, and *Drechslera halodes* was restricted to petiole segments of Kovipathagai site. The occurrence of *Colletotrichum* sp. was found in petiole segments of Chengalpet site in addition to Guduvanchery.

A total of 120 isolates belonging to 13 species was isolated from rhizome. The endophytic fungal community of rhizome segments was dominated by *Curvularia lunata* (45%) followed by *Phoma* sp. (13.33%) and *Fusarium* sp. (13.33%). The occurrence of *Curvularia lunata* was high in rhizome segments from Ambattur, Chengalpet and Kovipathagai sites, while *Phoma* sp. was found

in rhizome segments of Chengalpet and Guduvanchery site and *Fusarium* sp. was found in rhizome segments of Chengalpet and Avadi sites.

Root segments yielded a mere 34 isolates belonging to 5 species which was comparatively the lowest. The endophytic population of roots was dominated by *Curvularia lunata* (52.94%) and Mycelia Sterile (29.41%). *Curvularia lunata* was found to occur in high frequency in Ambattur, Chengalpet and Kovipathagai. The occurrence of sterile forms was restricted to the root segments of Guduvanchery and Kovipathagai. The isolation frequency (IF %) was calculated for the sites studied and is tabulated in Table 1.




Fig. 1 – The colonization frequency of endophytic fungi from various plant parts

The colonization frequency of endophytic fungi from various plant parts is showed in fig. 1. It could be observed that the colonization percentage was comparatively low for root segments with maximum colonization percentage of 12% in Ambattur and Kovipathagai root segments. High colonization frequency of >45% in all petiole segments was observed. The highest frequency of 98% was observed with petiole segments from Avadi. Leaflet and Rhizome segments had moderate frequency of colonization.

Jaccard's species diversity:

The Jaccard's species diversity of the endophytic fungal diversity isolated from the plant *Marsilea minuta* is presented in Table 2. The data obtained was subjected to statistical analysis such as Gleason, Shannon and Simpson's Dominance index to evaluate the richness, evenness and dominance of the endophytic fungal species isolated. It was observed that the overall endophytic fungal community had Gleason, Shannon and Simpson's index value to be 2.67, 1.94 and 4.88 respectively. It was found that the overall endophytic fungal diversity had low evenness with the Pielou's evenness of 0.31. This is due to the occurrence of certain fungi in low numbers.

The similarity between species isolated from different plants parts of *Marsilea minuta* from four different sites in and around Chennai is presented in Table 3. It could be observed that plant parts of same site had high similarity of species. Among the sites, Avadi had high similarity of species within its plant parts and Chengalpet had comparatively low similarity of species within its plant parts. The root segments showed no similarity of species with other plant parts in many cases.

Table 1 Relative density of Colonization (rD %) of Endophytic fungi from four sites in and around Chennai

	Relative density of Colonization (%)															
	Ambattur				Chengalpet				Guduvanchery				Avadi			
	L	P	R	RT	L	P	R	RT	L	P	R	RT	L	P	R	RT
<i>Acremonium</i> sp.	0	0	0	0	7.14	0	25	0	0	0	0	0	0	0	0	0
<i>Alternaria alternata</i>	22.22	0	0	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Aureobasidium pullulans</i>	0	0	0	0	0	0	0	0	2.77	0	0	0	0	0	0	0
<i>Botryodiplodia theobramae</i>	11.11	0	0	0	0	23.33	0	0	0	0	0	0	0	0	0	0
<i>Colletotricum</i> sp.	0	0	5.26	0	0	10	0	0	94.44	85.71	0	0	0	0	0	0
<i>Curvularia clavata</i>	0	0	5.26	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Curvularia lunata</i>	22.22	95.83	63.15	83.33	85.71	36.66	25	50	0	0	0	0	30	24.48	50	50
<i>Curvularia tuberculata</i>	0	0	5.263	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Drechslera australiensis</i>	0	0	0	0	7.14	0	0	0	0	0	0	0	0	0	0	0
<i>Drechslera halodes</i>	0	0	0	0	0	0	0	0	0	0	0	0	65	36.73	17.85	0
<i>Fusarium</i> sp.	0	0	0	0	0	0	25	0	0	0	0	0	5	12.24	25	16.66
<i>Nigrospora sphaerica</i>	0	0	5.26	0	0	0	0	0	0	0	0	0	0	0	0	0
<i>Penicillium funiculosum</i>	11.11	0	5.26	0	0	0	0	50	0	0	0	0	0	0	0	0
<i>Penicillium</i> sp.	22.22	0	0	16.66	0	0	0	0	0	0	0	0	0	0	0	0
<i>Phoma</i> sp.	0	0	0	0	0	30	25	0	2.77	10.71	77.77	0	0	0	0	0
<i>Scolecobasidium humicola</i>	0	0	5.26	0	0	0	0	0	0	0	0	0	0	0	3.57	0
Sterile form	11.11	4.16	5.26	0	0	0	0	0	0	3.57	22.22	100	0	24.48	0	33.33
<i>Torula</i> sp.	0	0	0	0	0	0	0	0	0	0	0	0	0	2.04	3.57	0

Ambattur was represented by the highest number of endophytic fungal species (11 species) followed by Chengalpet with 8 species. Guduvanchery and Avadi were represented by 6 fungal species each. The number of isolates was high in Avadi (206 isolates) followed by Guduvanchery (152 isolates), Ambattur (116 isolates) and Chengalpet (100 isolates).

It could also be observed that low similarity of species was observed within plant parts of different regions.

Table 2 The statistical analysis of the richness, evenness and dominance of endophytic fungal community isolated from the plant parts of *Marsilea minuta*.

	Site	Ni	Np	H _G	H _S	H _{Si}
Leaflet	Ambattur	18	6	1.72	1.73	5.4
	Chengalpet	28	4	0.9	0.51	1.08
	Guduvanchery	72	3	0.46	0.25	2.39
	Avadi	40	3	0.54	0.79	1.38
Petiole	Ambattur	48	2	0.25	0.17	1.34
	Chengalpet	60	4	0.73	1.29	3.46
	Guduvanchery	56	3	0.49	0.49	4
	Avadi	98	5	0.87	1.39	2
Rhizome	Ambattur	38	8	1.92	1.37	1.11
	Chengalpet	8	4	1.44	1.38	1.33
	Guduvanchery	18	2	0.34	0.52	1.52
	Avadi	56	5	0.99	1.23	1
Root	Ambattur	4	2	0.72	0.69	1.94
	Chengalpet	12	2	0.40	0.45	3.69
	Guduvanchery	6	1	0	0	2.88
	Avadi	12	3	0.80	1.01	2.57
All segments	Overall	574	18	2.67	1.94	4.88

Where *Ni*-Number of isolates; *Np*-Number of species; *H_G*-Gleason index; *H_S*-Shannon index; *H_{Si}*-Simpson's dominance index

Table 3 Jaccard's Similarity index between species isolated from different plant parts of *Marsilea minuta* from four different sites in and around Chennai

	L-A	P-A	R-A	Ro-A	L-C	P-C	R-C	Ro-C	L-G	P-G	R-G	Ro-G	L-Av	P-Av	R-Av	Ro-Av
L-A	1															
P-A	0.33	1														
R-A	0.27	0.25	1													
Ro-A	0.33	0.33	0.11	1												
L-C	0.12	0.25	0.1	0	1											
P-C	0.12	0.2	0.2	0	0.16	1										
R-C	0.11	0.2	0.09	0	0.2	0.33	1									
Ro-C	0.14	0.2	0.25	0	0.25	0.2	0.2	1								
L-G	0	0	0.12	0	0	0.4	0.2	0	1							
P-G	0.12	0.25	0.22	0	0	0.4	0.16	0	0.66	1						
R-G	0.14	0.33	0.11	0	0	0.2	0.2	0	0.33	0.66	1					
Ro-G	0.16	0.5	0.12	0	0	0	0	0	0	0.33	0.5	1				
L-Av	0.12	0.25	0.1	0.25	0.25	0.16	0.4	0.25	0	0	0	0	1			
P-Av	0.22	0.4	0.18	0.16	0.16	0.12	0.28	0.16	0	0.14	0.16	0.2	0.6	1		
R-Av	0.1	0.2	0.18	0.16	0.16	0.12	0.28	0.16	0	0	0	0	0.6	0.66	1	
Ro-Av	0.28	0.66	0.22	0.25	0.25	0.16	0.4	0.16	0	0.2	0.25	0.33	0.5	0.6	0.33	1

L- Leaflet; P- Petiole; R-Rhizome; Ro-Root; A-Ambattur; C-Chengalpet; G- Guduvanchery; Av-Avadi

Discussion

Studies on the endophytic community associated with aquatic Pteridophytes are scarce. The endophytic diversity from plant parts of *Marsilea minuta* collected from various sites were evaluated in the present study. The present study is one of its kind in reporting the endophytic fungal diversity from *Marsilea minuta*, an aquatic Pteridophyte. Altogether 17 species was identified which include the sterile mycelia.

Differences in endophytic fungal species was observed in the plant collected from different sampling sites. Some species were found abundant in particular region, while few species were restricted to a particular sampling site. It was observed that *Colletotrichum* sp. which was dominant in Guduvanchery, was meagre in other sites, *Curvularia lunata*, the most dominant fungi recorded in this study was completely absent in Guduvanchery. Collado et al. (1999), Gallery et al (2007) have reported such variations occurring due to differences in geographical location of the host. The comparison of diversity between two regions has been drawn by calculating the similarity index between the populations (Danilov & Ekelund 1999, Washington 1984). The statistical analysis showed that the diversity of the endophytic community varied greatly between various sites in the present study. The reason for variation in number of species and their diversity according to the site of collection of the plant has been elaborately explained in previous literatures (Petrini et al. 1992, Sieber 2007).

The fungal species which were isolated as endophytes exhibited tissue specificity and differences in endophytic fungal community among various plant parts. The occurrence of *Scolecobacidium humicola*, *Curvularia tuberculata*, *C. clavata* and *Nigrospora sphaerica* were restricted to petiole segments. While leaflet segments was the only host for *Drechslera australiensis*, *Aureobasidium pullulans* and *Alternaria alternata*. These observations indicate that the fungal species have affinity to specific tissue of the host and are able to utilize or survive within the substrate (Photita et al. 2001, Rodrigues 1994). It was observed that petiole segments harbored more number of colonies followed by leaflets, rhizome and root. This is due the proximity of petiole segments to the soil. The presence of endophytes in leaves are due to the exposure. Specificity in host plant part by endophytic fungi has been reported by Ganley & Newcombe (2006) and Huang et al. (2008).

The statistical analysis of the data provided thorough insight of the variation in diversity. The richness and evenness were statistically evaluated to indicate the plant parts or regions rich in particular species. Gleason index is sensitive to the richness of the fungal community. High Gleason index observed in leaflet and petiole of Ambattur and Rhizome of Chengalpet refers to low richness of the fungal species. Shannon index is a measure of both richness and evenness. Higher values describes that the species are equally distributed within the sample while values closer to 0 indicate that the species are not equally distributed within the population (Jost 2006). High values of Shannon index were observed in many parts of the plant indicating that the endophytic fungal community were rich and even. It was observed that the Shannon index for the overall plant was higher than plant parts, which might be due to the fact that no other plant part has complete diversity of all endophytic fungal species.

The root segments from Guduvanchery had Shannon index value of 0, which indicates dominance of single species (Sterile form). Leaflet segments of Guduvanchery and petiole segments of Ambattur had low Shannon index as they were dominated by *Colletotrichum* sp. and *Curvularia lunata* respectively. Simpson's dominance index which indicates the evenness of a fungal species, was found to be high in petiole of Guduvanchery and leaflet of Ambattur. The results indicate that these segments had an even distribution of fungal species.

Only three Coelomycetes namely, *Phoma* sp., *Colletotrichum* sp., and *Botryodiplodia thoebraeae* were isolated in the present study. Though in general the endophytic population is dominated by the Coelomycetes, their diversity was low in the present study. The possible reason for the host plant failing to harbour these species is the environmental stress which may have hindered the colonisation of Coelomycetes, as the sampling sites are largely polluted regions. This variation may be partly due to environmental conditions which include humidity, temperature,

rainfall and inoculum conditions (Photita et al. 2001, Santamaria & Bayman 2005). The study deals with four different geographical locations. However, expansion of the study from different aquatic plants of different nature will result in more diverse fungal species.

References

- Arnold AE, Lutzoni F. 2007 – Diversity and host range of foliar fungal endophytes, are tropical leaves biodiversity hotspots? *Ecology* 8, 541–549.
- Arnold AE, Maynard Z, Gilbert GS, Coley PD, Kursar TA. 2000 – Are tropical fungal endophytes hyperdiverse? *Ecology Letters* 3, 267–274.
- Azevedo JL, Araujo WL. 2007 – Diversity and applications of endophytic fungi isolated from tropical plants. In: Ganguli BN, Deshmukh SK (eds), *Multifaceted microbes*. CRC press, Boca Raton, 189–207.
- Bills GF. 1996 – Isolation and analysis of endophytic fungal communities from woody plants. In: *Endophytic Fungi in Grasses and Woody Plants* (eds. S.C. Redlin, L.M. Carris and M.N. St Paul). APS Press, USA, 31–65.
- Bhattamisra SK, Singh PN, Singh SK, Kumar V. 2007 – Anxiolytic activity of *Marsilea minuta* Linn in rodents. *Journal of Herbal Medicine and Toxicology* 1, 15–20.
- Carroll G. 1991 – Beyond pest deterrence – alternative strategies and hidden costs of endophytic mutualisms in vascular plants. In: Andrews J, Hirano SS, eds. *Microbial Ecology of Leaves*. New York: Springer Verlag, 358–375.
- Collado J, Platas G, Gonzales I, Pelaez F. 1999 – Geographical and seasonal influence on the distribution of fungal endophytes in *Quercus ilex*. *New Phytology* 143, 525–532.
- Danilov RA, Ekelund NGA. 1999 – The efficiency of seven diversity and one similarity indices based on phytoplankton data for assessing the level of eutrophication in lakes of central Sweden. *The Science of the Total Environment* 234, 15–23.
- Dobranic JK, Johnson IA, Alikhan QR. 1995 – Isolation of endophytic fungi from eastern larch (*Larix laricina*) leaves from New Brunswick, Canada'. *Canadian Journal of Microbiology* 41, 194–198.
- Ellis MB. 1971 – *Dematiaceous Hyphomycetes*. Commonwealth Mycological Institute, Kew. England. 608 p.
- Ellis MB. 1976 – *More Dematiaceous Hyphomycetes*. Commonwealth Mycological Institute, Kew. England. 507 p.
- Fisher PJ, Petrini O. 1987 – Location of fungal endophytes in tissue of *Suaeda fruticosa*: A preliminary study. *Transactions of the British Mycological Society* 89, 246–249.
- Frohlich J, Hyde KD, Petrini O. 2000 – Endophytic fungi associated with palms. *Mycological Research* 104, 1202–1212.
- Gallery RA, Dalling JW, Arnold AE. 2007 – Diversity, host affinity and 15 distribution of seed-infecting fungi: A case study with *Cecropia*. *Ecology* 88, 582–588.
- Ganley RJ, Newcombe G. 2006 – Fungal endophytes in seeds and needles of *Pinus monticola*. *Mycological Research* 110, 318–332.
- Hirsch G, and Braun U. 1992 – Communities of parasitic microfungi. In: Winterhoff W, ed. *Handbook of vegetative science, Volume 19. Fungi in vegetation science*. Dordrecht, the Netherlands: Kluwer Academic Publishers, 225–250.
- Huang WY, Cai YZ, Hyde KD, Corke H, Sun M. 2008 – Biodiversity of endophytic fungi associated with 29 traditional Chinese medicinal plants. *Fungal Diversity* 33, 61–75.
- Hyde KD, Soyong K. 2008 – The fungal endophyte dilemma. *Fungal Diversity* 33, 163–173.
- Jost L. 2006 – Entropy and diversity. *Oikos* 113, 363–375. doi: 10.1111/j.2006.0030-1299.14714.x
- Kumar DSS, Hyde KD. 2004 – Biodiversity and tissue-recurrence of endophytic fungi in *Tripterygium wilfordii*. *Fungal Diversity* 17, 69–90.

- Kumar DSS, Lau CS, Wan JMF, Yang D, Hyde KD. 2005 – Immunomodulatory compounds from *Pestalotiopsis leucothēs*. Fungal Diversity 87, (HKUCC 10197), an endophytic fungus of *Tripterygium wilfordii*. Life Sciences 78, 147–156.
- Li HY, Zhao CA, Liu CJ, Xu XF. 2010 – Endophytic fungi diversity of aquatic/riparian plants and their antifungal activity in vitro. Journal of Microbiology 1, 1–6.
- Madhu S, Kannabirran V, Frank PR, Reddy MS, Gnanasekar N. 2012 - Evaluation of anti diabetic activity of *Marsilea minuta* Linn against alloxan induced diabetes in albino rats. International Research Journal of Pharmacy 3, 223–225.
- Parihar P, Daswani L, Bohra A. 2003 – Toxic effect of plant of *Marsilea minuta* L. on the growth of *Staphylococcus aureus*. Indian Fern Journal 20, 48–50.
- Parrotta JA. 2001 – Healing medicinal plants of peninsular India. CAB International Publishing, New York, 24–25.
- Petrini O, Sieber TN, Toti L, Viret O. 1992 – Ecology, metabolite production and substrate utilization in endophytic fungi. Natural Toxins 1, 185–196.
- Photita W, Lumyong S, Lumyong P, Hyde KD. 2001 – Endophytic fungi of wild banana (*Musa acuminata*) at Doi Suthep Pui National Park, in Thailand. Mycological Research 105, 1508–1513.
- Qadri M, Rajput R, Abdin MZ, Vishwakarma RA, Riyaz-Ul-Hassan S. 2014 – Diversity, molecular phylogeny, and bioactive potential of fungal endophytes associated with the Himalayan blue pine (*Pinus wallichiana*). Microbial Ecology 67, 877–887.
- Rocha ACS, Garcia D, Uetanabaro APT, Carneiro RTO et al. 2011 – Foliar endophytic fungi from *Hevea brasiliensis* and their antagonism on *Microcyclus ulei*. Fungal Diversity 47, 75–84. <http://dx.doi.org/10.1007/s13225-010-0044-2>.
- Rodrigues KF. 1994 – The foliar endophytes of the Amazonian palm *Euterpe oleracea*. Mycologia 86, 376–385
- Sakalidis ML, Hardy GESJ, Burgess TI. 2011 – Endophytes as potential pathogens of the baobab species *Adansonia gregorii*: a focus on the Botryosphaeriaceae. Fungal Ecology 1, 1–14.
- Santamaria J, Bayman P. 2005 – Fungal epiphytes and endophytes of coffee leaves (*Coffea arabica*). Microbial Ecology 50, 1–8. <http://dx.doi.org/10.1007/s00248-004-0002-1>.
- Selim KA, El-Beih AA, AbdEl-Rahman TM, ElDiwany AI. 2011 - Biodiversity and antimicrobial activity of endophytes associated with Egyptian medicinal plants. Mycosphere, 2, 669–678.
- Sieber TN. 2007 – Endophytic fungi in forest trees: are they mutualists? Fungal Biol. Rev. 21, 75–89
- Stone JK, Polishook JD, White JRJ. 2004 – Endophytic fungi. In: Mueller G, Bills GF, Foster MS, eds. Biodiversity of fungi: inventory and monitoring methods. Burlington, MA, USA: Elsevier, 241–270.
- Suryanarayanan TS, Senthilarasu G, Muruganandam V. 2000 – Endophytic fungi from *Cuscuta reflexa* and its host plants. Fungal Diversity 4, 117–123.
- Suryanarayanan TS. 1992 – Light-incubation: a neglected procedure in mycology. The Mycologist 6, 144.
- Sutton BC. 1980 – The Coelomycetes. Commonwealth Mycological Institute, Kew, England. 696 p
- Tiwari S, Singh S, Pandey P, Saikia SK et al. 2014 – Isolation, structure determination, and anti-aging effects of 2,3-pentanediol from endophytic fungus of *Curcuma amada* and docking studies. Protoplasma DOI 10.1007/s00709-014-0617-0.
- Udaya Prakash NK. 2004 – Indoor molds: Isolation and Identification. Color Wings (M) Pvt. Ltd. Chennai, India, pp.99.
- U“Ren, Lutzoni F, Miadlikowska J, Laetsch AD, Elizabeth Arnold A. 2012 – Host and geographic structure of endophytic and endolichenic fungi at a continental scale. American Journal of Botany 99, 898–914.
- Washington HG. 1984 – Diversity, biotic and similarity indices. A review with special relevance to aquatic ecosystems. Water Research 18, 653–694.

White JF Jr, Torres MS. 2010 – Is plant endophyte-mediated defensive mutualism the result of oxidative stress protection? *Physiol Plant* 138(4), 440–6.